

REVIEW PAPER

## Selenium nanoparticles role in organ systems functionality and disorder

Seyed Mohammad Amini<sup>1,2\*</sup>, Vahid Pirhajati Mahabadi<sup>3</sup>

<sup>1</sup> Radiation Biology Research Center, Iran University of Medical Sciences, Tehran, Iran

<sup>2</sup> Medical Nanotechnology Department, School of Advanced Technologies in Medicine, Iran University of Medical Sciences, Tehran, Iran

<sup>3</sup> Neuroscience Research Center, Iran University of Medical Sciences, Tehran, Iran

### ARTICLE INFO

#### Article History:

Received 2 April 2018

Accepted 06 June 2018

Published 01 July 2018

#### Keywords:

Antioxidant

Nanopharmacokinetic

Nanotoxicity

Organ Function

Selenium Nanoparticles

### ABSTRACT

Extensive research on the nutritional and medical application of selenium nanoparticles (SeNPs) was performed in past decades. Besides nutritional values, new characteristics such as antibacterial and anticancer properties depict a bright future for high Selenium (Se) consumption in the coming years. Se is essential for the proper functioning of most of the major body organ systems meanwhile it could be highly toxic and even cancerous. The current knowledge of Se interaction with major organ systems functionality such as the central nervous system isn't well studied and many physiological aspects aren't clear to the science community. Meanwhile, various results were published on increasing organ system functionality through administrated SeNPs. So with the rapid entrance of SeNPs in the medical and nutritional industry, it may cause unintended complications. The intent of this review is to investigate current knowledge of SeNPs interaction with major body organ systems functionality. Investigated pharmacokinetic parameters of SeNPs was also reviewed.

### How to cite this article

Amini SM, Pirhajati Mahabadi V. Selenium nanoparticles role in organ systems functionality and disorder. *Nanomed Res J*, 2018; 3(3): 117-124. DOI: 10.22034/nmrj.2018.03.001

## INTRODUCTION

Se is an essential mineral with a high nutritional value and very important role for majority of physiological interactions in our body [1]. Our body intake of selenium is provided from food or water as a selenite, selenate, selenocysteine and selenomethionine and was carried by different human selenoproteins in the body. Humans with Selenium deficiency are amenable to many Health problems in a way that abnormal selenoprotein function or selenium deficiency could lead to various neurological [2], thyroid [1], muscle [3] and many other physiological disorders [4]. So the need for selenium starts from an embryonic age and continues until old age.

Nanoparticles are gaining lots of attention and concerns in biomedical and industrial

application [5, 6]. Especially metal nanoparticles with extraordinary characteristics are capable of many diagnostic [7, 8], therapeutic [9, 10], health [11, 12] and nutrition [13] application. Recently many reports about a different biomedical application of Se nanoparticles were published. With antimicrobial, antioxidant and anticancer properties [14] and lower toxicity in comparison to the selenite ( $\text{SeO}_4^{-2}$ ) or selenite ( $\text{SeO}_3^{-2}$ ) counterpart [15, 16], these nanoparticles have attracted much attention.

As a direct result of higher bioavailability of Se in nanoparticles form in comparison to other Se product, there is a growing need for Se nanoparticles in the livestock industry. Trial studies on animal feeding represents higher body weight gain [17], fertility rate [18], rumen fermentation [19], feed efficiency [20], and antioxidant status [21].

\* Corresponding Author Email: [Amini.sm@iums.ac.ir](mailto:Amini.sm@iums.ac.ir)  
[Mohammadamini86@gmail.com](mailto:Mohammadamini86@gmail.com)

Similar to livestock, agriculture industry applied Se nanoparticles for better plants growth and productivity [22]. Furthermore, the extensive use of Se as an antimicrobial agents in hygiene products was recommended by many researchers [23, 24].

Future extensive use of Se nanoparticles may lead to possible harmful effects on human health and the environment. Therefore the present review provides possible pros and cons of Se nanoparticle role in human organ functionality and disorder.

#### *Importance of Se for organ functionality*

The discovery of a congestive cardiomyopathy known as Keshan disease and Kashin-Beck disease in certain part of the People's Republic of China, which is deforming arthritis reviled the important role of Se in dietary [25]. But Keshan disease is not the only disorder of Se deficiency. Altogether, many experimental bioassays, epidemiologic studies, and even human clinical intervention trials support that a tiny amount of Se is vital for proper function of many human organs and systems such as Reproductive system [26], Thyroid [27], brain [28], cardiovascular system [29], Jacobson's organ function [30] and many others.

There is much evidence that indicates the direct relation between Se deficiency and increasing the risk of cancer [31, 32]. Molecular scientific bases of Se intake and cancer prevention are not clear but Glutathione peroxidase activity for hydrogen peroxidase breakdown strongly depends on available selenium [33]. Also, mitochondrial electron transport was altered in Se deficiency conditions [34] that could lead to more oxidative stress [35].

Malnutrition may also be associated with microbial infections. In case of Se deficiency, this has been reported frequently by researchers. Different research groups demonstrate that harmless viruses could be virulent in Se-deficient hosts [36, 37]. Boyne *et al.* reported that Se deficiency may have a negative effect on the mice immune system to eliminate *Candida albicans* [38]. Same results were reported for mice infected with *Diplococcus pneumoniae* [39] and *Listeria monocytogenes* [40]. Also, Viral infection leads to high ROS generation [41] and in-fact viral gene expression controls the cellular gene expression via ROS generation which leads to the appearance of cancer cells [42]. It's clear that under low Se condition, general oxidative stress increased and the mentioned process lead to higher viral infection. Also, Se deficiency may

cause a decrease in immune system efficiency that provides a chance for opportunistic pathogens [43, 44].

#### *Se toxicity and organ disorder*

Se toxicity was well-known in livestock that grazed in high Se soils. These animals developed disorders such as "alkai disease" and "blind staggers" [45]. Alkai disease is characterized by thickening hair and degenerative changes in hooves and blind staggers is characterized by impairment of vision and unsteady steps. In human, acute selenium toxicity usually leads to gastrointestinal disorders such as nausea, vomiting or diarrhoea alongside hair losses and headache [46]. Other acute toxicological symptoms such as bronchitis [47] and hypochromic anemia [48] have been reported. Long-term Se toxicity resembles in acute discoloration of skin and nail [49]. Except for the blood level of selenium, other biochemical indicators of body Se would increase which include: prothrombin time [50], alanine serum concentrations [51] and alanine aminotransferase enzyme activity [52]. Liver histopathological studies reveal sinusoidal damage and nodular regenerative hyperplasia in a rat model with high Se intake [53].

Numerous experimental studies demonstrate the carcinogenic effect of inorganic or organic Se form [54, 55] which is contrary to the cancer-protective activity of Se that was discussed. The same contradictive reports happen to the neuroprotective or neurodegenerative activity of Se. Amyotrophic lateral sclerosis (ALS) is a rare neurodegenerative disease that may be related to high Se intake [56]. Other nervous system abnormality may also be related to Se toxicity [57, 58]. Also, the limited evidence demonstrates Se toxicity in endocrine [48], reproductive [59], immune [60] systems. A direct link between high Se exposure and dental caries is clarified by Hadjimarkos *et al* [61].

#### *Pharmacokinetic parameters of Se NPs*

The science of investigating the scale and rate of absorption, distribution, metabolism, and elimination of drugs in the body through meticulous experimental methods known as a pharmacokinetic [61]. Pharmacokinetic values defined by applying specific parameters include the volume of distribution (Vd), half-life ( $T_{1/2}$ ), mean residence time (MRT), clearance (Cl), maximum concentration ( $C_{max}$ ), bioavailability (F) and area

under the time-concentration curve (AUC) (Fig. 1). In general metallic nanoparticles pharmacokinetic is different from inorganic or organic metallic ion forms [62].

In the human metabolic cycle, Se exists in various organic or inorganic oxidation states including -2, +2, +4 and +6 [63]. Unlike mentioned oxidative states of Se, pharmacokinetic data of elemental Se ( $Se^0$ ) is scarce. In nature, some microorganisms reduced selenite ( $Se^{+4}$ ), or selenate ( $Se^{+6}$ ) into nanoparticles formed  $Se^0$  [64]. Moderate absorption of  $Se^0$ NPs reduced by bacteria in the chicken model was reported [65]. Bioavailability and bioactivity of chemically synthesized  $Se^0$ NPs are reported in Rats [66]. The retention of SeNPs and Se-methylselenocysteine (MeSeCys) at the nutritional level in mice blood and other tissues was compared and no significant differences were observed [67]. Similar results were observed for kidney and liver accumulation for  $Se^0$ NP and selenite [68]. Same results were reported for selenomethionine (SeMet) [69] and selenite [68]. The clearance of SeNPs from the body can be either by renal and hepato-biliary excretion SeNPs or excretion of Se metabolites. In a pharmacokinetic study by Loeschner *et al.* the entrance of  $Se^0$  from

$Se^0$ NPs into the metabolic cycle and its subsequent elimination was confirmed by the apprehension of the two urinary metabolites of Se alongside a high amount of stool excretion of Se [68]. The authors don't investigate hair excretion.

#### Studies that indicate organ functionality with Se Nanoparticles Treatments

Organ functionality strongly depends on Antioxidant defensive capability. The antioxidant defence system is consisting of enzymatic and non-enzymatic antioxidants which are expanded not only inside the cells but also in extracellular environments. The first line of the antioxidant defence system is to restrain the production of free radicals. Reduction of disintegrated hydrogen peroxide and hydroperoxides by water and alcohol is a prime ROS generation mechanism which was suppressed through some selenoproteins and selenoenzymes such as glutathione peroxidase, thioredoxin reductases and etc. In many studies that applied SeNPs, increasing selenoproteins and selenoenzymes activity was reported with lower toxicity in comparison to an organic or inorganic form of Se [62]. SeNPs provides not only higher bioavailability but also they can scavenge free




Fig. 1: Pharmacokinetic values

radicals directly [63]. The organ systems in which the effect of selenium nanoparticles on their function is examined is shown schematically in Fig. 2.

**Musculoskeletal system:** Beside direct interaction with oxidative stress SeNPs could be integrated with other metabolic cycles that are involved with the musculoskeletal system oxidative stress response. For example, increasing oxidative stress can lead to a higher expression of heat shock proteins (HSPs) [64], which is an adjusting process corresponding to the interruption of cellular integrity [65]. The nutritional supplement of SeNPs could increase the expression level of HSP90 in a rest period after extensive exercise. A higher level of HSP expression could increase cells tolerability in stressful condition [66]. The nutritional supplement of SeNPs can significantly decrease the blood urea nitrogen (BUN) and creatinine serum concentration after intense exercise [67]. Both of these compounds can damage cells through oxidative stress [68].

**Reproductive system:** The main cause of sperm dysfunction is oxidative stress [18]. Some researchers applied SeNPs as a male fertility preservation substance. Increase semen quality and reproductive functionality of male goats was

reported by Shi *et al.* through increase activity of testicular GPx and improvement in testicular microstructure and testicular spermatozoa. Moreover, SeNPs treatment leads to higher Se accumulation in testes, and testicular tissue [19]. Fertility preservation in those who have been treated with chemotherapy is major concerns. Rezvanfar *et al.* observed the chemoprotective activity of SeNPs against cisplatin-induced gonadotoxicity. They demonstrate significant improvement in spermatogenesis and semen quality and lessened spermatid DNA destruction alongside cisplatin-induced free radical stress [69].

**Endocrine system:** Se is necessary for efficient endocrine glands functionality. Many selenoenzymes and selenoproteins are adjusting redox status of endocrine cells and in some cases involved in endocrine hormone metabolism [70]. But few investigations were performed for investigation of SeNPs role in endocrine glands functionality. Hassanin *et al.* demonstrated that accumulated SeNPs in thyroid cell can modify thyroid chromium toxicity in rats [71]. Rezaeian and Sadeghi report investigate the thyroid and sex hormones levels after SeNPs administration in rats. SeNPs administration can increase both thyroids and sex hormones levels not only in




Fig. 2: Schematic representation of SeNPs possible interaction with body organ systems: In this manuscript, current knowledge about SeNPs interaction with musculoskeletal, reproductive, endocrine, nervous, immune, digestive, circulation and the integumentary system was discussed.

normal condition but also in oxidative stress condition [72]. Oral administration of SeNPs can not only minimize diabetic complication but also could enhance pancreatic efficiency. Al-Quraishy *et al* report not only a significant augment of blood serum insulin level was observed but also histopathological studies show a SeNPs protective role in  $\beta$ -cells in the islet of Langerhans [73]. Similar results were provided by a liposomal formulation of SeNPs by Ahmed *et al* [74].

**Nervous system:** For normal nervous system function selenoproteins are essential. Nervous system damage due to nanoparticle was reported before [75]. A link between selenoproteins functionality and several nervous system disorder was reviewed by Schweizer *et al* [28]. But SeNPs were merely applied for investigation of central nervous system functionality. Naziroğlu *et al.* review state that SeNPs could be a potential remedial compound for Alzheimer disease treatment [76]. In contrast, Yuan *et al.* studied the impacts of SeNPs on sodium currents on dorsal root ganglion (DRG) neurons activity, applying the whole-cell patch clamp technique. In their report, SeNPs were able to decrease sodium current in a concentration and time-dependent manners which consider possible neurotoxic characteristic for SeNPs [77].

**Immune system:** It has been demonstrated that Se is generally accumulated in immune response organs such as liver, lymph nodes and spleen [78]. To deal with immunogens, Se treatment can increase antibodies production and expand complement responses through different mechanisms which include various selenoproteins [78]. It has been demonstrated that biogenic SeNPs could be a stimulator for immune system [80]. In particular, expanded neutrophils chemotactic and respiratory burst activities were observed for SeNPs in comparison with sodium selenite [81]. The authors consider the different pharmacokinetics parameters of SeNPs in comparison to selenite for assessment of immune stimulator functionality of SeNPs.

**Digestive system:** Sarkar *et al.* claimed that SeNPs could be an effective treatment for fatty liver disease [82]. In a separate study induced fatty liver male rats were treated with SeNP, and lower level of free radical and inflammation were reported [83]. SeNPs can improve rumen fermentation in livestock. A diet which contains SeNPs could increase total fiber digestion in sheep

[19]. *Lactobacillus* species could be applied for the preparation of selenium nanoparticle-enriched probiotics that represent an antifungal activity against *Candida albicans* [84].

**Circulation system:** Selenoproteins and selenoenzymes are essential for blood cells protections against oxidative damages [85]. It has been demonstrated that feeding of SeNPs has a better impact in comparison to sodium selenite upon peroxidative damage in blood cells [21]. SeNPs is also representing a better iron homeostasis capability in comparison to other Se forms. In vivo comparison investigation between SeNPs and sodium selenite, the level of transferrin and transferrin receptor was monitored. Total iron binding capacity (TIBC) in groups that treated by SeNPs is higher than a group that treated with sodium selenite [85]. Se has an essential role in inhibiting cardiovascular problems by augmenting the oxidation stress defence system to combat the oxidative alteration of lipids and subsequent decrease platelets aggregation [29]. SeNPs Ability to protect the cardiac cells from ischemia was demonstrated in an in-vitro experiment by Soumya *et al* [87].

**Integumentary system:** Se compensates skin-damaging compounds destructive role. It has been used for skin and hair care through cosmetics and health products. It has been demonstrated that SeNPs are able to penetrate in skin tissue and subsequently prevent the lipofuscin formation caused by UV-exposure and protect glutathione peroxidase (GPx) activity in mice [88]. Anti-fungal effect of SeNPs was as well as classical antidandruff selenium sulfide with lower cytotoxicity [89].

## CONCLUSION

As described throughout this review, because of the extreme role of Se in many selenoproteins and selenoenzymes function, a great deal of research is required to enhance our knowledge of SeNPs interaction in organ functionality and possible role in unwanted disorders. Digestive and integumentary absorption routes of SeNPs is not clearly identified but promising in-vitro and in-vivo results was reported for different organ systems functionality. SeNPs not only represent lower toxicity and higher antimicrobial and antioxidant properties but also it can be used in a larger nutritional amount. Sometimes, some technological profits could be potential problems. Large-scale addition of SeNPs as a livestock food

supply could happen very soon and subsequently human or environmental safety concerns such as long-term toxic and possible carcinogenic effects must be considered through further studies.

### CONFLICTS OF INTEREST

The authors declare that there are no conflicts of interest regarding the publication of this manuscript.

### REFERENCES

- Ventura M, Melo M, Carrilho F. Selenium and Thyroid Disease: From Pathophysiology to Treatment. *International Journal of Endocrinology*. 2017;2017:1-9.
- Pillai R, Uyehara-Lock JH, Bellinger FP. Selenium and selenoprotein function in brain disorders. *IUBMB Life*. 2014;66(4):229-39.
- Lescure A, Rederstorff M, Krol A, Guicheney P, Allamand V. Selenoprotein function and muscle disease. *Biochimica et Biophysica Acta (BBA) - General Subjects*. 2009;1790(11):1569-74.
- Schweizer U, Dehina N, Schomburg L. Disorders of selenium metabolism and selenoprotein function. *Current Opinion in Pediatrics*. 2011;23(4):429-35.
- Zarchi AAK, Amini SM, Farsangi ZJ, Mohammadi E, Moosavi Z, Harati PG. A study on the possibility of drug delivery approach through ultrasonic sensitive nanocarriers. *Nanomedicine Journal*, 2018;5 (3):127-137.
- Shirkhanloo H, Safari M, Amini SM, Rashidi M. Novel Semisolid Design Based on Bismuth Oxide (Bi<sub>2</sub>O<sub>3</sub>) nanoparticles for radiation protection. *Nanomedicine Research Journal*, 2017;2 (4):230-238.
- Emami T, Madani R, Golchinfar F, Shoushtary A, Amini SM. Comparison of Gold Nanoparticle Conjugated Secondary Antibody with Non-Gold Secondary Antibody in an ELISA Kit Model. *Monoclonal Antibodies in Immunodiagnosis and Immunotherapy*. 2015;34(5):366-70.
- Fatemi F, Amini SM, Kharrazi S, Rasaei MJ, Mazlomi MA, Asadi-Ghalehni M, et al. Construction of genetically engineered M13K07 helper phage for simultaneous phage display of gold binding peptide I and nuclear matrix protein 22 ScFv antibody. *Colloids and Surfaces B: Biointerfaces*. 2017;159:770-80.
- Amini SM, Kharrazi S, Jaafari MR. Radio frequency hyperthermia of cancerous cells with gold nanoclusters: an in vitro investigation. *Gold Bulletin*. 2017;50(1):43-50.
- Karimi Zarchi AA, Amini SM, Salimi A, Kharazi S. Synthesis and characterisation of liposomal doxorubicin with loaded gold nanoparticles. *IET Nanobiotechnology*. 2018;12(6):846-9.
- Darabpour E, Kashef N, Amini SM, Kharrazi S, Djavid GE. Fast and effective photodynamic inactivation of 4-day-old biofilm of methicillin-resistant *Staphylococcus aureus* using methylene blue-conjugated gold nanoparticles. *Journal of Drug Delivery Science and Technology*. 2017;37:134-40.
- Shaabani E, Amini SM, Kharrazi S, Tajerian R. Curcumin coated gold nanoparticles: synthesis, characterization, cytotoxicity, antioxidant activity and its comparison with citrate coated gold nanoparticles. *Nanomedicine Journal*, 2017;4 (2):115-125.
- Amini SM, Gilaki M, Karchani M. Safety of nanotechnology in food industries. *Electronic physician*, 2014;6 (4):962.
- Hosnedlova B, Kepinska M, Skalickova S, Fernandez C, Ruttkay-Nedecky B, Peng Q, et al. Nano-selenium and its nanomedicine applications: a critical review. *International Journal of Nanomedicine*. 2018;Volume 13:2107-28.
- Benko I, Nagy G, Tanczos B, Ungvari E, Sztrik A, Eszenyi P, et al. Subacute toxicity of nano-selenium compared to other selenium species in mice. *Environmental Toxicology and Chemistry*. 2012;31(12):2812-20.
- Shakibaie M, Shahverdi AR, Faramarzi MA, Hassanzadeh GR, Rahimi HR, Sabzevari O. Acute and subacute toxicity of novel biogenic selenium nanoparticles in mice. *Pharmaceutical Biology*. 2012;51(1):58-63.
- Shi L-g, Yang R-j, Yue W-b, Xun W-j, Zhang C-x, Ren Y-s, et al. Effect of elemental nano-selenium on semen quality, glutathione peroxidase activity, and testis ultrastructure in male Boer goats. *Animal Reproduction Science*. 2010;118(2-4):248-54.
- Aitken RJ, Baker MA. Oxidative stress and male reproductive biology. *Reproduction, Fertility and development*, 2004;16 (5):581-588.
- Shi L, Xun W, Yue W, Zhang C, Ren Y, Liu Q, et al. Effect of elemental nano-selenium on feed digestibility, rumen fermentation, and purine derivatives in sheep. *Animal Feed Science and Technology*. 2011;163(2-4):136-42.
- Hu CH, Li YL, Xiong L, Zhang HM, Song J, Xia MS. Comparative effects of nano elemental selenium and sodium selenite on selenium retention in broiler chickens. *Animal Feed Science and Technology*. 2012;177(3-4):204-10.
- Sadeghian S, Kojouri GA, Mohebbi A. Nanoparticles of Selenium as Species with Stronger Physiological Effects in Sheep in Comparison with Sodium Selenite. *Biological Trace Element Research*. 2011;146(3):302-8.
- El-Ramady H, Abdalla N, Taha HS, Alshaal T, El-Henawy A, Faizy SEDA, et al. Selenium and nano-selenium in plant nutrition. *Environmental Chemistry Letters*. 2015;14(1):123-47.
- Huang X, Chen X, Chen Q, Yu Q, Sun D, Liu J. Investigation of functional selenium nanoparticles as potent antimicrobial agents against superbugs. *Acta Biomaterialia*. 2016;30:397-407.
- Tran PA, Webster TJ. Antimicrobial selenium nanoparticle coatings on polymeric medical devices. *Nanotechnology*. 2013;24(15):155101.
- Chen X, Yang G, Chen J, Chen X, Wen Z, Ge K. Studies on the relations of selenium and Keshan disease. *Biological Trace Element Research*. 1980;2(2):91-107.
- Barrington JW, Lindsay P, James D, Smith S, Roberts A. Selenium deficiency and miscarriage: a possible link? *BJOG: An International Journal of Obstetrics and Gynaecology*. 1996;103(2):130-2.
- Olivieri O, Girelli D, Azzini M, Stanzial AM, Russo C, Ferroni M, et al. Low Selenium Status in the Elderly Influences Thyroid Hormones. *Clinical Science*. 1995;89(6):637-42.
- Schweizer U, Bräuer AU, Köhrle J, Nitsch R, Savaskan NE. Selenium and brain function: a poorly recognized liaison. *Brain Research Reviews*. 2004;45(3):164-78.
- N??ve J. Selenium as a risk factor for cardiovascular diseases. *Journal of Cardiovascular Risk*. 1996;3(1):42-7.
- Berliner DL, Monti-Bloch L, Jennings-White C, Diaz-Sanchez V. The functionality of the human vomeronasal

- organ (VNO): Evidence for steroid receptors. *The Journal of Steroid Biochemistry and Molecular Biology*. 1996;58(3):259-65.
31. Willett W, Steven Morris J, Pressel S, Taylor J, Frank Polk B, Stampfer M, et al. PREDIAGNOSTIC SERUM SELENIUM AND RISK OF CANCER. *The Lancet*. 1983;322(8342):130-4.
  32. Shamberger RJ. Relationship of selenium to cancer. I. Inhibitory effect of selenium on carcinogenesis. *Journal of the National Cancer Institute*, 1970;44 (4):931-936.
  33. Lawrence RA, Burk RF. Glutathione peroxidase activity in selenium-deficient rat liver. *Biochemical and Biophysical Research Communications*. 1976;71(4):952-8.
  34. Rani P, Lalitha K. Evidence for altered structure and impaired mitochondrial electron transport function in selenium deficiency. *Biological Trace Element Research*. 1996;51(3):225-34.
  35. Liu Y, Fiskum G, Schubert D. Generation of reactive oxygen species by the mitochondrial electron transport chain. *Journal of Neurochemistry*. 2002;80(5):780-7.
  36. Beck MA, Shi Q, Morris VC, Levander OA. Rapid genomic evolution of a non-virulent Coxsackievirus B3 in selenium-deficient mice results in selection of identical virulent isolates. *Nature Medicine*. 1995;1(5):433-6.
  37. Beck MA, Levander OA, Handy J. Selenium Deficiency and Viral Infection. *The Journal of Nutrition*. 2003;133(5):1463S-7S.
  38. Boyne R, Arthur JR. The Response of Selenium-Deficient Mice to *Candida albicans* Infection. *The Journal of Nutrition*. 1986;116(5):816-22.
  39. Hoffmann PR, Berry MJ. The influence of selenium on immune responses. *Molecular Nutrition & Food Research*. 2008;52(11):1273-80.
  40. Wang C, Wang H, Luo J, Hu Y, Wei L, Duan M, et al. Selenium deficiency impairs host innate immune response and induces susceptibility to *Listeria monocytogenes* infection. *BMC Immunology*. 2009;10(1):55.
  41. Love AJ. Cauliflower mosaic virus, a Compatible Pathogen of Arabidopsis, Engages Three Distinct Defense-Signaling Pathways and Activates Rapid Systemic Generation of Reactive Oxygen Species. *PLANT PHYSIOLOGY*. 2005;139(2):935-48.
  42. Waris G, Ahsan H. Reactive oxygen species: role in the development of cancer and various chronic conditions. *Journal of carcinogenesis*, 2006;5:14.
  43. Descamps-Latscha B, Jungers P, Witko-Sarsat V. Immune System Dysregulation in Uremia: Role of Oxidative Stress. *Blood Purification*. 2002;20(5):481-4.
  44. Laddha NC, Dwivedi M, Mansuri MS, Gani AR, Ansarullah M, Ramachandran AV, et al. Vitiligo: interplay between oxidative stress and immune system. *Experimental Dermatology*. 2013;22(4):245-50.
  45. Kim YY, Mahan DC. Biological Aspects of Selenium in Farm Animals. *Asian-Australasian Journal of Animal Sciences*. 2003;16(3):435-44.
  46. Schuh B, Jappe U. Selenium intoxication: undesirable effect of a fasting cure. *British Journal of Dermatology*. 2007;156(1):177-8.
  47. Nemery B. Metal toxicity and the respiratory tract. *European Respiratory Journal*, 1990;3 (2):202-219.
  48. Nagai I. An Experimental Study of Selenium Poisoning. *Igaku Kenkyu*, 1959;29 (5):1505-1532.
  49. Smith MI, Westfall BB. Further Field Studies on the Selenium Problem in Relation to Public Health. *Public Health Reports (1896-1970)*. 1937;52(40):1375.
  50. Yang G, Yin S, Zhou R, Gu L, Yan B, Liu Y. Studies of safe maximal daily dietary Se-intake in a seleniferous area in China. Part II: Relation between Se-intake and the manifestation of clinical signs and certain biochemical alterations in blood and urine. *Journal of trace elements and electrolytes in health and disease*, 1989;3 (3):123-130.
  51. Longnecker MP, Taylor PR, Levander OA, Howe M, Veillon C, McAdam PA, et al. Selenium in diet, blood, and toenails in relation to human health in a seleniferous area. *The American Journal of Clinical Nutrition*. 1991;53(5):1288-94.
  52. Coudray C, Haida H, Boucher F, Tirard V, De Leiris J, Favier A. Effect of selenium supplementation on biological constants and antioxidant status in rats. *Journal of Trace Elements in Medicine and Biology*. 1996;10(1):12-9.
  53. Dubuisson L, Boussarie L, Bedin C-A, Balabaud C, Bioulac-Sage P. Transformation of sinusoids into capillaries in a rat model of selenium-induced nodular regenerative hyperplasia: An immunolight and immunoelectron microscopic study. *Hepatology*. 1995;21(3):805-14.
  54. Birt DF, Pour PM, Pelling JC. The Influence of Dietary Selenium on Colon, Pancreas, and Skin Tumorigenesis. *Selenium in Biology and Medicine: Springer Berlin Heidelberg*; 1989. p. 297-304.
  55. Woutersen RA, Appel MJ, van Garderen-Hoetmer A. Modulation of Pancreatic Carcinogenesis by Antioxidants. *Food and Chemical Toxicology*. 1999;37(9-10):981-4.
  56. Vinceti M, Guidetti D, Pinotti M, Rovesti S, Merlin M, Vescovi L, et al. Amyotrophic Lateral Sclerosis after Long-Term Exposure to Drinking Water with High Selenium Content. *Epidemiology*. 1996;7(5):529-32.
  57. Pamphlett R, McQuilty R, Zarkos K. Blood Levels of Toxic and Essential Metals in Motor Neuron Disease. *NeuroToxicology*. 2001;22(3):401-10.
  58. Yang GQ, Wang SZ, Zhou RH, Sun SZ. Endemic selenium intoxication of humans in China. *The American Journal of Clinical Nutrition*. 1983;37(5):872-81.
  59. Vinceti M, Cann CI, Calzolari E, Vivoli R, Garavelli L, Bergomi M. Reproductive outcomes in a population exposed long-term to inorganic selenium via drinking water. *Science of The Total Environment*. 2000;250(1-3):1-7.
  60. Ferenčík M, Ebringer L. Modulatory effects of selenium and zinc on the immune system. *Folia Microbiologica*. 2003;48(3):417-26.
  61. Hadjimarkos TIDM. Selenium in Relation to Dental Caries. *Food and Cosmetics Toxicology*. 1973;11(4):1083-95.
  62. Zhang J, Wang X, Xu T. Elemental Selenium at Nano Size (Nano-Se) as a Potential Chemopreventive Agent with Reduced Risk of Selenium Toxicity: Comparison with Se-Methylselenocysteine in Mice. *Toxicological Sciences*. 2007;101(1):22-31.
  63. Peng D, Zhang J, Liu Q, Taylor EW. Size effect of elemental selenium nanoparticles (Nano-Se) at supranutritional levels on selenium accumulation and glutathione S-transferase activity. *Journal of Inorganic Biochemistry*. 2007;101(10):1457-63.
  64. Khassaf M, McArdle A, Esanu C, Vasilaki A, McArdle F, Griffiths RD, et al. Effect of Vitamin C Supplements on Antioxidant Defence and Stress Proteins in Human Lymphocytes and Skeletal Muscle. *The Journal of*

- Physiology. 2003;549(2):645-52.
65. Kinnunen S, Hyypää S, Oksala N, Laaksonen DE, Hannila ML, Sen CK, et al.  $\alpha$ -Lipoic acid supplementation enhances heat shock protein production and decreases post exercise lactic acid concentrations in exercised standardbred trotters. *Research in Veterinary Science*. 2009;87(3):462-7.
  66. Kregel KC. Invited Review: Heat shock proteins: modifying factors in physiological stress responses and acquired thermotolerance. *Journal of Applied Physiology*. 2002;92(5):2177-86.
  67. Kojouri GA, Sharifi S. Preventing Effects of Nano-Selenium Particles on Serum Concentration of Blood Urea Nitrogen, Creatinine, and Total Protein During Intense Exercise in Donkey. *Journal of Equine Veterinary Science*. 2013;33(8):597-600.
  68. Zhang Z, Dmitrieva NI, Park JH, Levine RL, Burg MB. From The Cover: High urea and NaCl carbonylate proteins in renal cells in culture and in vivo, and high urea causes 8-oxoguanine lesions in their DNA. *Proceedings of the National Academy of Sciences*. 2004;101(25):9491-6.
  69. Rezvanfar MA, Rezvanfar MA, Shahverdi AR, Ahmadi A, Baeeri M, Mohammadirad A, et al. Protection of cisplatin-induced spermatotoxicity, DNA damage and chromatin abnormality by selenium nano-particles. *Toxicology and Applied Pharmacology*. 2013;266(3):356-65.
  70. Beckett GJ, Arthur JR. Selenium and endocrine systems. *Journal of Endocrinology*. 2005;184(3):455-65.
  71. Hashem K, Hassnin KMA, Abdel-Kawi SH. The prospective protective effect of selenium nanoparticles against chromium-induced oxidative and cellular damage in rat thyroid. *International Journal of Nanomedicine*. 2013;1713.
  72. Rezaeian-Tabrizi M, Sadeghi AA. Plasma antioxidant capacity, sexual and thyroid hormones levels, sperm quantity and quality parameters in stressed male rats received nano-particle of selenium. *Asian Pacific Journal of Reproduction*. 2017;6(1):29-34.
  73. Abdel Moneim A, Al-Quraishy S, Dkhil MA. Anti-hyperglycemic activity of selenium nanoparticles in streptozotocin-induced diabetic rats. *International Journal of Nanomedicine*. 2015;6741.
  74. Ahmed HH, Abd El-Maksoud MD, Abdel Moneim AE, Aglan HA. Pre-Clinical Study for the Antidiabetic Potential of Selenium Nanoparticles. *Biological Trace Element Research*. 2016;177(2):267-80.
  75. Amiri S, Yousefi-Ahmadipour A, Hosseini M-J, Haj-Mirzaian A, Momeny M, Hosseini-Chegeni H, et al. Maternal exposure to silver nanoparticles are associated with behavioral abnormalities in adulthood: Role of mitochondria and innate immunity in developmental toxicity. *NeuroToxicology*. 2018;66:66-77.
  76. Naziroğlu M, Muhamad S, Pecze L. Nanoparticles as potential clinical therapeutic agents in Alzheimer's disease: focus on selenium nanoparticles. *Expert Review of Clinical Pharmacology*. 2017;10(7):773-82.
  77. Yuan H, Lin J, Lan T. Effects of Nano Red Elemental Selenium on Sodium Currents in Rat Dorsal Root Ganglion Neurons. 2006 International Conference of the IEEE Engineering in Medicine and Biology Society; 2006/08: IEEE; 2006.
  78. Dickson RC, Tomlinson RH. Selenium in blood and human tissues. *Clinica Chimica Acta*. 1967;16(2):311-21.
  79. McKenzie RC, Rafferty TS, Beckett GJ, Arthur JR. Effects of selenium on immunity and aging. *Selenium: Springer US*; 2001. p. 257-72.
  80. Yazdi M, Mahdavi M, Setayesh N, Esfandyar M, Shahverdi A. Selenium nanoparticle-enriched *Lactobacillus brevis* causes more efficient immune responses in vivo and reduces the liver metastasis in metastatic form of mouse breast cancer. *DARU Journal of Pharmaceutical Sciences*. 2013;21(1):33.
  81. Kojouri GA, Sadeghian S, Mohebbi A, Mokhber Dezfouli MR. The Effects of Oral Consumption of Selenium Nanoparticles on Chemotactic and Respiratory Burst Activities of Neutrophils in Comparison with Sodium Selenite in Sheep. *Biological Trace Element Research*. 2011;146(2):160-6.
  82. Sarkar B, Bhattacharjee S, Daware A, Tribedi P, Krishnani KK, Minhas PS. Selenium Nanoparticles for Stress-Resilient Fish and Livestock. *Nanoscale Research Letters*. 2015;10(1).
  83. Hegedüs V, Prokisch J, Fébel H, Kleiner D, Ditrói K, Szijártó A, et al. Nanoselenium treatment in fatty liver. *Zeitschrift für Gastroenterologie*. 2012;50(05).
  84. Kheradmand E, Rafii F, Yazdi M, Sepahi A, Shahverdi A, Oveisi M. The antimicrobial effects of selenium nanoparticle-enriched probiotics and their fermented broth against *Candida albicans*. *DARU Journal of Pharmaceutical Sciences*. 2014;22(1):48.
  85. Murthy NLN. Investigation Of Biochemical And Histopathological Modifications Induced By Selenium Toxicity On Albino Mice. 2013.
  86. Kojouri GA, Jahanabadi S, Shakibaie M, Ahadi AM, Shahverdi AR. Effect of selenium supplementation with sodium selenite and selenium nanoparticles on iron homeostasis and transferrin gene expression in sheep: A preliminary study. *Research in Veterinary Science*. 2012;93(1):275-8.
  87. Soumya RS, Vineetha VP, Salin Raj P, Raghu KG. Beneficial properties of selenium incorporated guar gum nanoparticles against ischemia/reperfusion in cardiomyoblasts (H9c2). *Metallomics*. 2014;6(11):2134-47.
  88. Zhai X, Zhang C, Zhao G, Stoll S, Ren F, Leng X. Antioxidant capacities of the selenium nanoparticles stabilized by chitosan. *Journal of Nanobiotechnology*. 2017;15(1).
  89. Parsamehr N, Rezaie S, Khodavaisy S, Salari S, Hadizadeh S, Kord M, et al. Effect of biogenic selenium nanoparticles on ERG11 and CDR1 gene expression in both fluconazole-resistant and -susceptible *Candida albicans* isolates. *Current Medical Mycology*. 2017;3(3):16-20.